

Name: _____ Sex: ___ B.D.: _____ Pt#: _____
SSN: _____ Phone (day): _____ (eve): _____
Address (required): _____

CHARGE TO: (please attach admissions face sheet, completed insurance forms, and/or a copy of insurance card. If not available, please complete.)
Name: _____ Phone: _____
Address: _____
Policy Holder: _____ ID#: _____ Group #: _____

SPECIMEN INFORMATION

Amniotic Fluid	Solid Tissue	Peripheral Blood
Chorionic Villi	Placenta	Other: _____
Weeks Gestation: (LMP) _____	Fetal Tissue	
(US) _____	Products of Conception	

Clinical Indication/ICD-10: _____
Advanced Maternal Age
Abnormal Maternal Serum Screen
Abnormal NIPT
Increased risk for Down Syndrome
Increased risk for _____
Ultrasound Abnormalities: _____
Recurrent Miscarriage
Family History of: _____
Other: _____

TESTING

Chromosome analysis
Limited Follow-Up Study (PB Only)
FISH: Aneuvysion (Chromosomes 13, 18, 21, X, Y)
POC Aneuploidy (Chromosomes 13, 16, 18, 21, 22, X, Y)
Other: _____
Chromosomal Microarray*
Report all findings
Do not report variants of uncertain significance
Molecular/DNA Testing*: _____
Culture cells for possible sendout testing
AFP* ACHE* Reflex to ACHE if AFP Positive*
Viral testing* for: _____
Other: _____

REFLEX TESTING

If Aneuvysion FISH is NORMAL, then reflex to chromosomal microarray.
 ABNORMAL, then reflex to chromosome analysis.
If _____ is NORMAL,
 ABNORMAL, then reflex to _____

Physician: _____ Address: _____
Time/Date: _____
Phone: _____
Fax: _____

SPECIMEN REQUIREMENTS AND SHIPPING

All specimens must be labeled with patient's name and be accompanied by completed request form. All samples should be kept at room temperature and transported to the laboratory with minimum delay. Please call (800) 328-2026 if you have any questions.

AMNIOTIC FLUID

Amniotic Fluid: (AF) 15-20 ml of amniotic fluid in 2-3 sterile tubes. Please use B-D syringe. Do not use: Monoject or Jelco syringes, or silicon coated tubes. AFP testing will be performed on all specimens unless otherwise indicated.

Interphase FISH: Rapid aneuploid screening on uncultured cells (AF or CVS) is performed only in conjunction with conventional metaphase chromosome analysis, no irreversible therapeutic action should be initiated on the basis of FISH. Aspirate 5 ml of amniotic fluid in addition to the 15 ml of fluid required for cytogenetics. Bloody samples are not appropriate for interphase FISH due to the increased risk of maternal cell contamination.

PERIPHERAL BLOOD

Peripheral Blood: 5-10 ml of blood in a preservative free sodium-heparin (green-top) tube. Invert tube to mix. Prometaphase analysis will be performed on all specimens unless otherwise specified.

Newborns & Percutaneous Umbilical Blood (PUBS): Minimum of 1 ml peripheral blood in a preservative free sodium-heparin (green top) tube. Invert tube to mix well.

SOLID TISSUE

All solid tissue samples should be collected aseptically and transported in tissue culture media or Hank's balanced salt solution. Do NOT put in water, fixative, formalin or saline. Please keep sample at room temperature.

Products Of Conception/Fetal Tissue: Large chorionic villi sample (approximately 1-3 cm³) and a fetal tissue sample such as skin, lung, or pericardium. Please send multiple tissue types if possible. Label tube with tissue type or origin.

Skin Biopsy/Solid Tissue: 1-3 mm³ or more tissue. Label tube with tissue type or origin.

NEOPLASIA

Bone Marrow: Aspirate 1-2 ml bone marrow into a sterile syringe containing 0.1 ml preservative free sodium heparin invert syringe to mix and transfer to a 3 ml preservative free sodium-heparin (green-top) vacutainer tube.

Leukemic Peripheral Blood: Patient should have WBC of 15,000 or higher with approximately 10% circulating immature myeloid or lymphoid blast cells. Collect 5 ml of peripheral blood in a preservative free sodium-heparin (green top) vacutainer tube.

Solid Tumor Tissue: >1 cm³ representative tumor tissue collected under aseptic conditions and transported in sterile tissue culture media.

Lymph Node Biopsy: >5 mm³ tumor biopsy collected under aseptic conditions and transported in sterile tissue culture media.

MOLECULAR ANALYSIS/DNA TESTING

Peripheral Blood: 5-10 ml blood in an EDTA (lavender-top) tube for molecular testing, and 5-10 ml blood in a preservative free sodium-heparin (green-top) tube for cytogenetic studies. (Molecular studies will be forwarded to an outside laboratory).

Prenatal: 15-20 ml of amniotic fluid in 2 sterile tubes. Cytogenetic analysis will be performed, and amniocytes will be cultured to send to an outside laboratory for molecular studies.

FLUORESCENCE IN SITU HYBRIDIZATION (FISH)

FISH studies are indicated when classic cytogenetics alone cannot resolve an abnormality. Specimen collection is as described previously for the tissue to be studied.

SHIPPING INFORMATION

Sample should be securely packaged and sent at room temperature to:

Diagnostic Cytogenetics, 2292 West Commodore Way Ste 100 Seattle, WA 98199

We provide free shipping. To arrange pick up in the local Puget Sound area, please call (206) 328-2026 or (800) 328-2026. For overnight delivery service: Federal Express (800) 463-3339. Call us for our current account number. Please send specimens by Standard Overnight Service. Specimens sent on Friday MUST be marked with a "Saturday Delivery" sticker.

Please call the lab at (800) 328-2026 with the airbill number so that we may track your specimen.